arXiv:1211.6761v1 [math.KT] 28 Nov 2012

Twisted K-theory constructions in the case of a decomposable Dixmier-Douady class II: Topological and equivariant models

Antti J. Harju

Abstract

We study twisted K-theory on a product space $\mathbb{T} \times M$. The twisting is given by a cup product class which applies the 1-cohomology of $\mathbb{T}$ and 2-cohomology of M. In the case of a topological product, we give a concrete realization of the gerbe transition functions associated to the cup product characteristic class and for twisted K^1-theory elements. The nontriviality of this construction is proved. We also study equivariant twisted K-theory and gerbes in the case a smooth group action on a manifold. The twisting comes from a product class. By applying groupoid-Cech hypercohomology of an action groupoid we find explicit realization for the equivariant gerbes and twisted K^1-elements. Moreover, we develop a superconnection formalism in equivariant cohomology which is used to extract information of the twisted K-theory classes.

MSC: 19L50, 53C08, 22A22.

1 Introduction

In the first part of this series we gave a concrete realization for elements of a K^1-groups twisted by a gerbe on a product manifold $\mathbb{T} \times M$. The twisting class decomposes as $\alpha \smile \beta \in H^3(\mathbb{T} \times M)$, for $\alpha \in H^1(\mathbb{T})$ and $\beta \in H^2(M)$. We used Hamiltonian quantization to produce gerbes associated to these classes: quantization leads to a projective bundle of Hilbert spaces whose characteristic class is the three form part of the local K^1-index theory on the parameter space of a Dirac family. The K-theory classes twisted by these gerbes were given explicitly using quantum field theory in the same spirit Wess-Zumino-Witten model has been used in the construction of equivariant twisted K-theory classes on a Lie group [15].

A gerbe associated to the decomposable class has a simple topological realization. There is a bundle of Fock Hilbert spaces over $\mathbb{T} \times M$. The Fock space has a basis which can be decomposed into different charge subspaces labeled by $\mathbb{Z}$. We fix a line bundle λ associated to a nontrivial class in $H^2(M, \mathbb{Z})$. Then we make the Fock states of charge $k \in \mathbb{Z}$ topologically equivalent to $\lambda^{\otimes k}$ over the manifold M. Moreover, we make the translations over the circle $\mathbb{T}$ change the charge of each state. Consequently there are transition functions valued in $PU(\mathcal{H})$ associated to the creation and annihilation processes of the bundle of topological type λ over M. In this case, the Dixmier-Douady (DD) class is the product class.

In [11] we used techniques from differential geometry to study the characteristic class of the gerbes as well as twisted K-theory group elements. These techniques involve the local families index theorem and superconnection techniques. Here we raise the issue of how to extend this formalism to the topological framework: the setup itself is still well defined on the topological product space $\mathbb{T} \times X$. The main problems are to prove the nontriviality of the gerbe and to find the precise twisted K^1-theory class of the construction. The characteristic class of the gerbe is solved completely. The twisted K^1-group elements are proven to be nontrivial.

We return to the smooth setup in the study of equivariant gerbes and equivariant twisted K-theory on a smooth manifold $\mathbb{T} \times M$. Our strategy to construct gerbes is to replace the twisting line bundles in the Fock bundle model with equivariant line bundles on M. The G action on $\mathbb{T}$ is chosen to be trivial. Here we work under the hypothesis that there is a cover which is invariant under the group action and the components of the cover are contractible to a finite number of points.

We use the Lie groupoid formalism which is flexible enough to allow lots of generalizations of this type. In this case we can give explicit formulas for the equivariant $PU(\mathcal{H})$-transition functions as well as DD class for the gerbe in terms of cocycles in groupoid cohomology. The DD class of the gerbe is the cup product class $\alpha \smile \beta_G$ where α is a 1-cohomology class associated to the trivial G-module structure in $\mathbb{T}$ whereas β_G is a cocycle of a G-equivariant complex line bundle.

We define twisted K^1-group elements as certain sections of a bundle of self adjoint Fredholm operators associated to the gerbe which are invariant under the group action. As in [11] the Mayer-Vietoris sequence is used to study the structure of these groups. We also define an equivariant superconnection in the twisted equivariant K^1-group which is mapped to the quotient of the equivariant cohomology $H_G^{\mathrm{odd}}(\mathbb{T} \times M)$ by an ideal generated by the wedge product of the superconnection form and the connection form of the equivariant twisting line bundle λ. This allows us to compare different equivariant classes.

As an example of equivariant gerbes and equivariant twisted K-theory we consider the action of the circle group $\mathbb{T}$ on the sphere S^2 which rotates the sphere around its axis. There are two fixed points: the poles. Consequently, there are nontrivial $\mathbb{T}$-equivariant line bundles parametrized by irreducible representations of $\mathbb{T}$ on the northern and the southern hemisphere. Such a line bundle is used to construct a $\mathbb{T}$-equivariant gerbe. The ordinary K-theory group $K^1_{\mathbb{T}}(\mathbb{T} \times M)$ is a sum of representation rings of $\mathbb{T}$ constrained by the common virtual dimension. The twisted K^1-groups are defined similarly but the generators of the representation rings are truncated by the representations on the fibres of λ. Silimar phenomenon occurs in twisted equivariant K-theory on Lie groups which was studied by Freed-Hopkins-Teleman, [8], [9]. We apply the superconnection analysis to study these groups.

By experiments it is usual that twising in K-theory produce torsion subgroups if the ordinary K-theory groups are torsion free. For example, in [11] we observed this in the case of $\mathbb{T}^3$ and $\mathbb{T} \times S^2$. More examples can be found in [3]. Twisted K-theory of the simple Lie groups is all torsion [7]. Here we observe a phenomenon contrary to this one. The ordinary K^1-groups of $\mathbb{T} \times \mathbb{R}P^{2n}$ have torsion subgroups. However, after twisting by a product class in the 3-cohomology of $\mathbb{T} \times \mathbb{R}P^{2n}$ the torsion subgroups vanish. Our Fredholm operator model gives a satisfactory explanation of this. Furhermore, we observe that the superconnection defines an isomorphism of groups $K^1(\mathbb{T} \times \mathbb{R}P^{2n}, \alpha \smile \beta) \to H^{\mathrm{even}}_{\mathrm{dR}}(\mathbb{T} \times \mathbb{R}P^{2n}, \mathbb{Z})$ which in noninjective when defined in the ordinary K-theory.

Conventions. $\mathcal{H}$ denotes an infinite dimensional complex Hilbert space. The group $U(\mathcal{H})$ is equipped with the strong operator topology and it is contractible. The symbols $\underline{\mathbb{T}}$ and $\underline{PU}(\mathcal{H})$ denote the sheaves of $\mathbb{T}$ and $PU(\mathcal{H})$ valued functions. In chapter 2. when we consider topological spaces these functions are supposed to be continuous and smooth in the rest of the work.

Denote by $\mathbf{Fred}^{(0)}$ and $\mathbf{Fred}^{(1)}$ the usual classifying spaces of K-theory: the space of bounded Fredholm operators and the space of bounded self-adjoint Fredholm operators with both positive and negative essenstial spectrum, both equipped with the norm topology. In this work we are only interested in unbounded self-adjoint operators. Let Ψ denote the map $Q \mapsto Q(1+Q^2)^{-1/2}$. Then we set $\mathbf{Fred}^{(1)}_{\Psi} = \Psi^{-1}(\mathbf{Fred}^{(1)})$. The space $\mathbf{Fred}^{(1)}_{\Psi}$ equipped with a topology induced by Ψ is homotopy equivalent to $\mathbf{Fred}^{(1)}$, [12]. Therefore, we can use unbounded Fredholm operators to classify (twisted) K-theory. If Y is a compact topological space and $\sigma \in H^3(Y, \mathbb{Z})$, it defines a $PU(\mathcal{H})$-bundle and there is an associated bundle of $\mathbf{Fred}^{(i)}$ operators, $i = 0, 1$. The action of $PU(\mathcal{H})$ on $\mathbf{Fred}^{(i)}$ is by conjugation. Then $K^i(Y, \sigma)$ is by definition the space of homotopy classes of sections of the Fredholm bundle. In the case $i = 1$ we can replace $\mathbf{Fred}^{(1)}$ by $\mathbf{Fred}^{(1)}_{\Psi}$. In the groupoid context in chapter 3 and beyond we introduce generalized twisted K-theory groups.

2 Twisted K-Theory of Topological Products $\mathbb{T} \times X$

2.1. Let $\mathcal{H}$ be an infinite dimensional complex Hilbert space with a polarization $\mathcal{H} = \mathcal{H}^+ \oplus \mathcal{H}^-$. A canonical anti commutation relations algebra, CAR, is a unital C^*-algebra generated by $a(f) : f \in \mathcal{H}$ such that $a : \mathcal{H} \to \mathrm{CAR}$ is an antilinear map and CAR is subject to

$$\{a(u), a(v)\} = 0 \quad \text{and} \quad \{a(u), a(v)^*\} = \langle u, v \rangle \mathbf{1},$$

for all $u, v \in \mathcal{H}$. A Fock space $\mathcal{F}$ is a Hilbert space with a vacuum vector $|0\rangle$ and CAR acts on the vacuum by $a(u)|0\rangle = 0 = a^*(v)|0\rangle$ for $u \in \mathcal{H}^+, v \in \mathcal{H}^-$ and such that the basis of $\mathcal{F}$ consists of

$$a(u_{i_1}) \cdots a(u_{i_k}) a^*(u_{j_1}) \cdots a^*(u_{j_l})|0\rangle \qquad \text{for} \qquad u_{i_\eta} \in \mathcal{H}^-, u_{j_\eta} \in \mathcal{H}^+. \tag{1}$$

This representation of CAR is irreducible. The basis vectors of this type are eigenvectors for the densely defined charge operator N, the eigenvalue of (1) being $l - k$. Then we can write the Fock space as a completion of the charge subspaces $\mathcal{F} = \widehat{\bigoplus}_{k \in \mathbb{Z}} \mathcal{F}^{(k)}$. The operator S denotes the shift operator which sends $\mathcal{F}^{(k)}$ to $\mathcal{F}^{(k+1)}$ for all $k \in \mathbb{Z}$. We refer to [18] for a careful introduction of these matters.

The Hilbert space vectors are labeled so that $\mathcal{H}^+$ has the basis $\{u_i : i \geq 0\}$ and $\mathcal{H}^-$ has the basis $\{u_i : i < 0\}$.

Consider a compact topological space X and a nontrivial complex line bundle λ on X. Let $\{V_i : i = 1 \leq i \leq n\}$ be a cover of X which trivializes λ with respect to the transition functions $h_{ij} : V_{ij} \to \mathbb{T}$. For $\mathbb{T}$ we choose a cover $\{\mathbb{T}_+, \mathbb{T}_-\}$ such that $\mathbb{T}_+ \cap \mathbb{T}_-$ consist of two disconnected arcs: neighborhoods of -1 and 1. We denote these subsets by $\mathbb{T}_{+-}^{(\pm 1)}$. Then we set an open cover $\{U_i : 1 \leq i \leq 2n\}$ for $\mathbb{T} \times X$ such that

$$U_i = \mathbb{T}_+ \times V_i \quad \text{and} \quad U_{i+n} = \mathbb{T}_- \times V_i \quad \text{for all} \quad 1 \leq i \leq n. \tag{2}$$

We will construct a projective Fock bundle over $\mathbb{T} \times X$ with a typical fibre $\mathcal{PF} = \mathcal{F}/\mathbb{T}$. We introduce transformations which raise (lower) the charges of the Fock states by one whenever we go around the circle in the positive (negative) direction. This can be achieved as follows. We define the local families $S : U_{i,j+n} \to PU(\mathcal{H})$ which raise the charge of the state. These operations are defined to be independent of $(\phi, p) \in U_{i,j+n}$. They are valued in $PU(\mathcal{H})$ since restricted to each fibre, S creates a state from the vacuum:

$$S|0\rangle = a^*(u_0)|0\rangle,$$

but we need to choose an orthonormal basis vector u_0 from the subspace it spans. These operators act on the CAR algebra under conjugations by

$$S a(u_k) S^{-1} = a(u_{k+1})$$

for all $k \in \mathbb{Z}$ and similarly for $a^*(u_k)$. We can fix a basis vector u_0 and introduce lifted local families $\widehat{S} : U_{i,j+n} \to U(\mathcal{H})$.

We make the charge k states in the lifted Fock bundle to be of topological type $\lambda^{\otimes k}$ over X. Thus, we define the cocycle in $g \in H^1(\mathbb{T} \times M, \underline{PU}(\mathcal{H}))$ by

$$\begin{aligned}
g_{ij}(\phi, p) &= (h_{ij}(p))^N, \\
g_{i+n,j+n}(\phi, p) &= (h_{i+n,j+n}(p))^N, \\
g_{i,j+n}^{(-1)}(\phi, p) &= (h_{i,j+n}(p))^N \\
g_{j+n,i}^{(-1)}(\phi, p) &= (h_{j+n,i}(p))^N \\
g_{i,j+n}^{(1)}(\phi, p) &= (h_{i,j+n}(p))^N S \\
g_{j+n,i}^{(1)}(\phi, p) &= S^{-1}(h_{j+n,i}(p))^N
\end{aligned}$$

The upper index (± 1) means that the cocycle is defined in the neighborhood of ± 1 in $\mathbb{T}$. The canonical lift of this cocycle has exactly the properties discussed above. We denote by **PF** the projective Fock bundle determined by g.

For the lifted functions, the gerbe cocycle relation reads

$$f_{abc}(\phi, p) = \widehat{g}_{ab}(\phi, p)\widehat{g}_{bc}(\phi, p)\widehat{g}_{ac}(\phi, p)^{-1}.$$

for all $1 \leq a, b, c \leq 2n$. In [11] we found out that the Dixmier-Douady cocycle is equal to the totally antisymmetric cocycle whose nonidentity components are determined by

$$f_{i,j+n,k+n}^{(1)}(\phi, p) = (h_{jk}(p))^{-1} \tag{3}$$

for $1 \leq i, j, k \leq n$. This is a consequence of the relation $SN = (N-1)S$ applied to the lifted transition functions.

Proposition. The Dixmier-Douady class of $\mathbf{PF}$ is equal to the cup product $\sigma := \alpha \smile \beta$ where α is a generator of the group $H^1(\mathbb{T})$ and β is the cohomology class of the twisting line bundle λ.

Proof. This can be proven by following the lines of the Proposition in 3.5. below in the case of a trivial group $G = \{1\}$. $\square$

We introduce another dependence on the topology of X by introducing an arbitrary rank complex vector bundle ξ over X and then define $\mathbf{PF}_\xi$ to be the projective bundle $\mathbf{P}(\mathbf{F} \otimes \xi)$. Without loss of generality, we can assume that both bundles λ and ξ are trivialized over U_i. Namely, we can always take refinements of the cover to make this happen. The shift operator S acts on this bundle by $S \otimes 1$. This does not have effect on the DD class.

2.2. Here we recall the construction of twisted K^1-groups from [11] which can be applied in the topological setup straightforwardly. We consider a real Clifford $*$-algebra generated by $\psi_n, n \in \mathbb{Z}$ and subject to

$$\{\psi_n, \psi_m\} = 2\delta_{n,-m}, \qquad \psi_n^* = \psi_{-n}.$$

We set a vacuum representation, $\mathcal{H}_s$, for the Clifford algebra such that ψ_0 acts on the vacuum by $\psi_0 \eta_0 = \eta_0$. The operators ψ_i with $i < 0$ annihilate the vacuum subspace whereas ψ_i with $i > 0$ are used to generate the basis of the module. Then we define trivial Hilbert bundles $U_i \times (\mathcal{H}_s \otimes \mathcal{F} \otimes \mathbb{C}^r)$ for each $1 \leq i \leq 2n$ where r is the rank of ξ. The local self-adjoint families $Q^i : U_i \to \mathbf{Fred}^{(1)}_\Psi(\mathcal{H}_s \otimes \mathcal{F} \otimes \mathbb{C}^r)$ are defined by

$$Q^i_{\phi,p} = (\sum_{n\in\mathbb{Z}} \psi_n \otimes e_{-n} + \frac{\phi}{2\pi}\psi_0 \otimes \mathbf{1}) \otimes \mathbf{1}.$$

We call Q a supercharge since restricted on $\mathbb{T}$ it defines a supercharge operator in a Wess-Zumino-Witten model. We have used the operators e_n which determine a representaion of the centrally extended Lie algebra of the loop group $L\mathbb{T}$ on $\mathcal{F}$ by

$$e_n = \sum_{i\in\mathbb{Z}} : a^*(u_{n+i}) a(u_i) : \qquad \text{with} \qquad [e_n, e_m] = -2n\delta_{n,-m}.$$

The usual normal ordering :: is applied to make these operators well defined in the Fock space, $: a^*(u_i)a(u_j) := -a(u_j)a^*(u_i)$ if $i = j < 0$ and the order is unchanged otherwise. Q is a family of unbounded operators. Then the $PU(\mathcal{H})$-valued cocycle g acts on the local families Q^i by conjugation according to

$$g_{i,j+n}(\phi, p) Q^{j+n}_{\phi,p} (g_{i,j+n}(\phi,p))^{-1} = Q^i_{\phi-2\pi,p}.$$

Therefore, Q^i are patched together by an adjoint action of the $PU(\mathcal{H})$ valued Cech 1-cocycle corresponding to the Dixmier-Douady class σ. Then it follows, by definition, that these families are group elements in $K^1(\mathbb{T} \times X, \sigma)$. The twisted K-theory group $K^1(\mathbb{T} \times X, \sigma)$ for a topological space X can be solved using the Mayer-Vietoris sequences as we did in [11], see theorem 2. The theorem can be used in the topological setup without modifications. Alternatively, the calculation in 3.6. does not require smoothness and can be applied here in the case of a trivial group $G = \{1\}$.

2.3. Next we study conditions for the nontriviality of a twisted K^1-group element. Let Y denote a compact topological space with nontrivial 3-cohomology. Let $\sigma \in H^3(Y, \mathbb{Z})$ which we realize as a projective Hilbert bundle $\mathbf{PH}_\sigma$ which trivializes over a finite cover $\{U_i\}$. Here Y and σ are arbitrary, not necessarily a product space or a product class as avove. Denote by g the transition functions of the projective bundle and by $\widehat{g}$ the lifted transition functions.

Consider $Q \in K^1(Y, \sigma)$ realized as local families of unbounded operators $Q^i : U_i \to \mathbf{Fred}^{(1)}_\Psi$ acting on local Hilbert bundles over U_i. Let us denote by $\mathcal{H}^{(a,b)}_{i,x}$ the (a,b)-spectral subspaces of Q^i_x

in the fibre $\mathcal{H}_{i,x}$ of the local Hilbert bundle over U_i. On $U_{ij} \neq \emptyset$ these spectral subspaces transform as $\widehat{g}_{ij}\mathcal{H}^{(a,b)}_{j,x} = \mathcal{H}^{(a,b)}_{i,x}$. However, there is no globally defined Hilbert bundle over Y. A σ-twisted spectral section P of Q is locally defined families of projection operators, $P^i : U_i \to \mathbf{Fred}^{(1)}_\Psi$ which transform equivariantly under the conjugation action of $\widehat{g}_{ij}$, that is, P is a section in the $\mathbf{Fred}^{(1)}_\Psi$-bundle associated to σ and for some positive real number $R > 0$ (depends on P) and every i and $x \in U_i$

$$\begin{aligned} P^i_x &= \mathbf{1} \quad \text{in} \quad \mathcal{H}^{(R,\infty)}_{i,x}, \\ P^i_x &= 0 \quad \text{in} \quad \mathcal{H}^{(-\infty,-R)}_{i,x}. \end{aligned} \tag{4}$$

The proof of the following theorem applies the ideas of the corresponding result in [14] in the ordinary K-theory.

Theorem. Suppose that the local families $Q^i : U_i \to \mathbf{Fred}^{(1)}_\Psi$ define a class in $K^1(Y,\sigma)$ and there is a neighborhood of zero $(-a,a) \subset \mathbb{R}$ so that the spectral subspaces $\mathcal{H}^{(-a,a)}_{i,x}$ are finite dimensional for all i and $x \in U_i$. Then Q has a spectral section if and only if Q is trivial in the group $K^1(Y,\sigma)$.

Proof. The condition $[Q] = 0$ in the twisted K^1-group realized by local families of $\mathbf{Fred}^{(1)}_\Psi$-valued sections is equivalent to $[Q(Q^2+1)^{-1/2}] = 0$ in the isomorphic twisted K^1-group realized by local families of $\mathbf{Fred}^{(1)}$-valued sections.

Assume that the spectral section exists. Then we define the local families $(Q')^i : U_i \to \mathbf{Fred}^{(1)}$

$$(Q')^i = \frac{Q^i}{\sqrt{(Q^i)^2+1}} - P^i \frac{Q^i}{\sqrt{(Q^i)^2+1}} P^i - (1-P^i)\frac{Q^i}{\sqrt{(Q^i)^2+1}}(1-P^i).$$

The operators $(Q')^i$ patch together to define a section in the σ twisted Fredholm bundle because P^i and Q^i do. For each i and $x \in U_i$, $(Q')^i_x$ restricts to a finite rank operator on the $(-R,R)$ spectral subspace and therefore Q' is a section of finite rank operators. Then, in norm topology we have the equality of homotopy classes $[Q(Q^2+1)^{-1/2} - Q'] = [Q(Q^2+1)^{-1/2}]$. The section $Q(Q^2+1)^{-1/2} - Q'$ is homotopic to the section Q'' which is locally defined by

$$(Q'')^i = P^i \frac{Q^i + r}{\sqrt{(Q^i)^2+1}} P^i + (1-P^i)\frac{Q^i - r}{\sqrt{(Q^i)^2+1}}(1-P^i)$$

for any $r \in \mathbb{R}$. For large enough r, the first term is strictly positive whereas the second term is strictly negative. Thus, there is an inverse section $(Q'')^{-1}$ of Q''. Then $[Q''] = 0$ and therefore $[Q(Q^2+1)^{-1/2}] = 0$.

We use the following lemmas to prove the converse statement.

Lemma [14]. Let A be a bounded self-adjoint operator on $\mathcal{H}$ such that $A^2 - A$ is of finite rank. Assume that there is a self-adjoint projection P such that $||A - P|| < 1/2$. Then $A : \mathrm{Im}(P) \to \mathrm{Im}(AP)$ is an isomorphism onto a closed subspace and if Π denotes the orthonormal projection onto $\mathrm{Im}(AP)$, then $A - \Pi$ has a finite rank.

Lemma. Let B be a section of a $\mathbf{Fred}^{(1)}$ bundle twisted by σ. If $U_{ij} \neq \emptyset$ and P^i and P^j are projections onto $\mathrm{Im}(B^i)$ and $\mathrm{Im}(B^j)$, then $\widehat{g}_{ij}P^j(\widehat{g}_{ij})^{-1} = P^i$ and thus P is a section.

Proof. Let $v \in \mathrm{Im}(B)$. Then $v_i = B^i w_i$ for some locally defined section w of the Hilbert bundle. If $U_{ij} \neq \emptyset$, then

$$\widehat{g}_{ij}P^j(\widehat{g}_{ij})^{-1}v_i = \widehat{g}_{ij}P^j(\widehat{g}_{ij})^{-1}B^i w_i = \widehat{g}_{ij}P^j B^j w_j = \widehat{g}_{ij}B^j w_j = B^i w_i = v_i.$$

Suppose then that $u_i(x) \notin \mathrm{Im}(B^i_x)$ for some $x \in U_i$. Then $P^i_x u_i(x) = 0$. If

$$\widehat{g}_{ij}(x)P^j_x(\widehat{g}_{ij}(x))^{-1}u_i(x) = \widehat{g}_{ij}P^j_x u_j(x) \neq 0.$$

Then $u_j(x) = B^j_x w_j(x) + w'_j(x)$ where $w_j(x) \neq 0$ and $w'_j(x) \notin \mathrm{Im}(B^j_x)$. Now

$$u_i(x) = \widehat{g}_{ij}(x)(B^j_x w_j(x) + w'_j(x)) = B^i(x)w_i(x) + w'_i(x)$$

where $w'_i(x) \notin \mathrm{Im}(B^i_x)$. Consequently $P^i_x u_i(x) \neq 0$. This is a contradiction.

Assume that Q is trivial in K-theory. Then we can set a homotopy $Q(t)$ between a locally constant section of invertible Fredholm operators, $Q(0)$, and $Q = Q(1)$ through sections in the σ twisted Fredholm bundle. We can set $Q(0)$ so that there are no eigenvalues in some arbitrarily small inverval $(0, \epsilon)$. Moreover, we choose the homotopy so that there are only finitely many eigenstates with eigenvalue in $(0, \epsilon)$ for any $Q^i_x(t)$ with $t \in [0, 1]$.

We define a continuous function $\chi \in C(\mathbb{R})$ such that $\chi(\lambda) = 0$ for $\lambda < 0$ and $\chi(\lambda) = 1$ for $\lambda \geq \epsilon$ and apply the spectral theorem to define

$$J^i(t) = \chi(Q^i(t)), \qquad t \in I := [0, 1].$$

Then $J(0)$ is a locally constant section of projection operators and $J(1)$ satisfies the condition (4) for $R = \epsilon$ but it may not be projection section, however $(J(1))^2 - J(1)$ is a finite rank section.

We divide the interval $[0, 1]$ to subintervals $[t', t'']$ on which $||J^i_x(t') - J^i_x(s)|| < 1/2$ for all $s \in [t', t'']$. We proceed inductively by proving that if the lower limit in the interval has a projection section, $P(t')$, such that $J(t') - P(t')$ is a finite rank section, then such a section exists for any t-value in this interval. At $t = 0$ the projection section exists. Consider $s \in [t', t'']$ and define a section $A^i_x(s) = J^i_x(s) + P^i_x(t') - J^i_x(t')$. These operators satisfy $||A^i_x(s) - P^i_x(t')|| < 1/2$ and $(A^i_x(s))^2 - A^i_x(s)$ is a finite rank operator for all i and $x \in U_i$. By the first lemma above, the orthogonal projection $P^i_x(s)$ onto the subspace $\mathrm{Im}(A^i_x(s)P^i_x(t'))$ in the fibre at x of the local Hilbert bundle over U_i differs from $A^i_x(s)$ and thus from $J^i_x(s)$ by a finite rank operator. By the second lemma, the projection families defined this way patch together to form a section. Thus, $t \mapsto P(t)$ defines a homotopy of projection sections and $J(t) - P(t)$ is a finite rank section for all $t \in [0, 1]$.

Consider the decomposition of the fibres of the local bundles $\mathcal{H}_i$ into the eigenspaces of Q^i. Then $J(1)$ is viewed as local families of diagonal operators. Next we define new local families $(P')^i$ by modifying $P^i(1)$ so that $J(1) - P'$ restricts to an operator which is nonzero in a finite span of eigenvectors of Q everywhere. This can be done as follows. Let $J(1) - P(1) = C$. Then we continuously truncate the eigenspace expansion of C to make it a section in the σ-twisted Fredholm bundle such that $(C')^i_x$ restricts to an operator on a finite span of Q^i eigenvectors for each $x \in X$. Let $(P')^i$ be the projection onto the image of

$$(P^i(1) + (C^i - (C')^i)P^i(1) = (J^i(1) - (C')^i)(J^i(1) - C^i)$$

for all i. The left side of this equation proves that $J^i(1) - (P')^i$ restricts to a finite rank operator everywhere. The image of the local section on the right side is easily seen to differ from the image of $J^i(1)$ by a finite span of eigenvectors of Q^i everywhere. The section P' is a twisted spectral section: we can pick R which is greater than any of the eigenvalues of Q in the image $J(1) - P'$, and greater than ϵ. $\square$

Corollary. The local families of supercharge operators on $\mathbb{T} \times X$ (introduced in 2.2.) define a nontrivial element in $K^1(\mathbb{T} \times X, \sigma)$.

Proof. The families $Q^i : U_i \to \mathbf{Fred}^{(1)}_\Psi$ do not have a spectral section because of the spectral flow around the unit circle. $\square$

3 Twisted K-theory of Action Groupoids

3.1. We return to the smooth setup: M is a compact smooth manifold. In addition we assume that there is a smooth action of a compact Lie group G on M and that there is a G-invariant cover $\{V_i : 1 \leq i \leq n\}$ of M such that the Cech cohomology groups $H^k(\coprod_i V_i, \mathbb{T})$ are equal to zero for all $k > 0$. This is the case, for example, if each component V_i is a contractible set. We follow the notation of 2.1. and set a cover $\{U_i : 1 \leq i \leq 2n\}$ for $\mathbb{T} \times M$. We apply a trivial G action on $\mathbb{T}$

and therefore, the cover $\{U_i\}$ is G-invariant and the Cech cohomology groups $H^k(\coprod_i U_i, \underline{\mathbb{T}})$ vanish for all $k > 0$.

We apply the theory of Lie groupoids to construct a G-equivariant gerbe. The action groupoid $G \times \mathbb{T} \times M \rightrightarrows \mathbb{T} \times M$ is Morita equivalent to the groupoid $\coprod_{ij} G \times U_{ij} \rightrightarrows \coprod_i U_i$. We return to this in 3.3. Then we can apply the groupoid cohomology to the latter groupoid and our analysis determines explicitly the roles of transformations between different components of the cover and group actions in the cohomology class of the gerbe.

3.2. Let $\Gamma \rightrightarrows X$ denote a compact Lie groupoid. For each $p \geq 1$ let $\Gamma^{(p)}$ denote the manifold of composable sequences of arrows. We also use $\Gamma^{(0)} = X$. We have $p+1$ maps

$$\partial_i : \Gamma^{(p)} \to \Gamma^{(p-1)} \quad \text{for} \quad 0 \leq i \leq p$$

such that ∂_0 leaves out the first arrow, ∂_p leaves out the last arrow and the rest of ∂_i compose the i'th and $i+1$'th arrow. Then we set $\partial : \Gamma^{(p)} \to \Gamma^{(p-1)}$ by $\partial = \sum_i (-1)^i \partial_i$.

Let F denote an abelian sheaf in the category of smooth manifolds and F^p a small sheaf induced by F on $\Gamma^{(p)}$. Then we choose injective resolutions $F^p \to I^{p\bullet}$. This defines a double complex $I^\bullet(\Gamma^\bullet)$ whose total cohomology groups, $H^k(\Gamma^\bullet, F)$, are the cohomology groups of the groupoids with values in the abelian sheaf F. This theory is Morita invariant [2], [20].

We can also use the de Rham complex in the case of coefficients in $\mathbb{R}$. The exterior derivative defines a map $d : \Lambda^k(\Gamma^{(p)}) \to \Lambda^{k+1}(\Gamma^{(p)})$. We can use this with the coboundary operator $\partial^* : \Lambda^k(\Gamma^{(p)}) \to \Lambda^k(\Gamma^{(p+1)})$ to construct a double complex. The total cohomology of this complex is the de Rham cohomology of $\Gamma^\bullet$, we write $H^k_{\mathrm{dR}}(\Gamma^\bullet) = H^k(\Lambda^\bullet(\Gamma^\bullet))$. The de Rham theorem is still valid in the form

$$H^k_{\mathrm{dR}}(\Gamma^\bullet) \simeq H^k(\Gamma^\bullet, \mathbb{R}).$$

We call a cocycle in de Rham theory an integral cocycle, if it maps under the isomorphism into the image of the canonical map $H^k(\Gamma^\bullet, \mathbb{Z}) \to H^k(\Gamma^\bullet, \mathbb{R})$.

We are interested in the cohomology groups $H^2(\Gamma^\bullet, \underline{\mathbb{T}})$ and $H^3(\Gamma^\bullet, \underline{\mathbb{Z}})$. We also point out the existence of the cohomology group $H^1(\Gamma^\bullet, \underline{PU}(\mathcal{H}))$. We only consider compact groupoids (and therefore proper) and consequently the group homomorphism $H^2(\Gamma^\bullet, \underline{\mathbb{T}}) \to H^3(\Gamma^\bullet, \underline{\mathbb{Z}})$, constructed from the standard exact sequence of groups

$$0 \to \mathbb{Z} \to \mathbb{R} \to \mathbb{T} \to 1,$$

is an isomorphism. The group homomorphism $H^1(\Gamma^\bullet, \underline{PU}(\mathcal{H})) \to H^2(\Gamma^\bullet, \underline{\mathbb{T}})$, constructed from the central extension $\mathbb{T} \to U(\mathcal{H}) \to PU(\mathcal{H})$, is not an isomorphism in general. However, there is a canonical left inverse, [20].

A $\mathbb{T}$-central extension of a groupoid $\Gamma \rightrightarrows X$ is a groupoid $\hat{\Gamma} \rightrightarrows X$ together with a groupoid morphism

$$(\pi, \mathrm{id}) : (\hat{\Gamma} \rightrightarrows X) \to (\Gamma \rightrightarrows X)$$

such that there is a left $\mathbb{T}$ action on $\hat{\Gamma}$ making $\pi : \hat{\Gamma} \to \Gamma$ a (left) principal $\mathbb{T}$-bundle. These two structures are required to satisfy the compatibibilty

$$(\mu \cdot x)(\mu' \cdot y) = \mu\mu' \cdot (xy)$$

for all $(x, y) \in \hat{\Gamma}^{(2)}$ and $\mu, \mu' \in \mathbb{T}$. In [20] the elements of $\mathrm{Ext}(\Gamma, \mathbb{T})$, the Morita equivalence classes of $\mathbb{T}$ extensions of Γ, were identified with isomorphism classes of $\mathbb{T}$-gerbes over the stack associated to Γ. The set $\mathrm{Ext}(\Gamma, \mathbb{T})$ can be equipped with a structure of an abelian group. Then the group $\mathrm{Ext}(\Gamma, \mathbb{T})$ is isomorphic to $H^2(\Gamma^\bullet, \underline{\mathbb{T}})$. We shall not need the notion of an gerbe over a stack or Morita equivalence in this work and we refer to [2], [20] for details. In our case, the twisting classes arise naturally from equivariant $PU(\mathcal{H})$-bundles. Therefore, we are given a $H^1(\Gamma^\bullet, \underline{PU}(\mathcal{H}))$ cocycle which determines a $H^2(\Gamma^\bullet, \underline{\mathbb{T}})$ cocycle. This is then related to the decomposable class $\alpha \smile \beta$ under the isomorphism onto $H^3(\Gamma^\bullet, \mathbb{Z})$.

In [20] the group of twistings of the K-theory is taken to be $H^2(\Gamma^\bullet, \underline{\mathbb{T}})$ or equivalently $H^3(\Gamma^\bullet, \mathbb{Z})$. Then the canonical left inverse is applied to get a cocycle in $H^1(\Gamma^\bullet, \underline{PU}(\mathcal{H}))$. As noted above, we don't need to worry about this ambiguity. Associated to a cocycle $g \in H^1(\Gamma^\bullet, \underline{PU}(\mathcal{H}))$ there is a principal $PU(\mathcal{H})$ bundle $P \to X$ over the groupoid $\Gamma \rightrightarrows X$. Then there is an associated bundle

$$\mathbf{Fred}^{(i)}(g) := P \times_{PU(\mathcal{H})} \mathbf{Fred}^{(i)} \to X$$

for $i = 0, 1$. The action of $PU(\mathcal{H})$ on $\mathbf{Fred}^{(i)}$ is by conjugation. We denote by $\Xi^{(i)}(g)$ the space of continuous Γ-invariant sections of the bundle $\mathbf{Fred}^{(i)}(g)$. The twisted K-theory is defined by

$$K^i(\Gamma^\bullet, g) = \{[Q] : Q \in \Xi^{(i)}(g)\} \tag{5}$$

for $i = 0, 1$ where $[Q]$ denotes the homotopy class of Q. We consider unbounded operators and therefore, we can replace the space $\mathbf{Fred}^{(1)}$ by a homotopy equivalent space $\mathbf{Fred}^{(1)}_\Psi$ in the definition (5).

3.3. Let G be a compact Lie group which acts smoothly on M. We work under the assumptions of 3.1. for the cover $\{V_i\}$ of M and $\{U_i\}$ of $\mathbb{T} \times M$.

Consider the Lie groupoid $\Gamma := \coprod_{ij} G \times U_{ij} \rightrightarrows \coprod_i U_i$ with $s(g, x_{ij}) = x_j$ and $t(g, x_{ij}) = gx_i$. The multiplication of composable pairs is defined by

$$(g, (hx)_{ij})(h, x_{jk}) = (gh, x_{ik}).$$

The groupoid Γ is Morita equivalent to the action groupoid $G \ltimes (\mathbb{T} \times M)$. Especially, the cohomology group $H^*(\Gamma^\bullet, \mathbb{Z})$ is isomorphic to the equivariant cohomology groups $H^*_G(\mathbb{T} \times M)$. Since M is compact, the equivariant cohomology can be computed with the Cartan or Weyl model. Equivalently, we can use the groupoid cohomology of the Morita equivalent groupoid Γ.

Let us fix a sheaf of abelian groups F. Recall that when X is a manifold, the sheaf cohomology of a compact space X associated to an injective resolution is isomorphic to a Cech cohomology computed from an open cover $\{N_i\}$ of X which is acyclic for the sheaf F. More precisely, the Cech cohomology groups $H^k(N_{i_1 \cdots i_j}, F)$ are trivial for all $k > 0$ and for any intersection $N_{i_1 \cdots i_j}$ of the components of $\{N_i\}$.

We can fix a good open cover $\{N_i^{(p)}\}$ for each manifold $\Gamma^{(p)}$ with $p > 0$. These covers are acyclic for any sheaf of abelian groups. We consider the sheaf $\underline{\mathbb{T}}$. The cover with only one open set $N^{(0)} = \Gamma^{(0)}$ makes the Cech resolution of $\Gamma^{(0)}$ acyclic. Then the groupoid double complex is equivalent to the Cech hypercohomology complex associated to the chosen resolution

$$\begin{tikzcd}
\cdots & \cdots & \cdots & \\
C^\infty(\Gamma^{(0)}, \mathbb{T}) \arrow[r] \arrow[u, "\delta"] & \prod_{ab} C^\infty(N^{(1)}_{ab}, \mathbb{T}) \arrow[r] \arrow[u, "\delta"] & \prod_{ab} C^\infty(N^{(2)}_{ab}, \mathbb{T}) \arrow[r] \arrow[u, "\delta"] & \cdots \\
C^\infty(\Gamma^{(0)}, \mathbb{T}) \arrow[r] \arrow[u, "\delta"] & \prod_{a} C^\infty(N^{(1)}_{a}, \mathbb{T}) \arrow[r] \arrow[u, "\delta"] & \prod_{a} C^\infty(N^{(2)}_{a}, \mathbb{T}) \arrow[r] \arrow[u, "\delta"] & \cdots
\end{tikzcd} \tag{6}$$

The horizontal lines are induced from the isomorphisms to the injective resolution and the choice is not unique. However, the cohomology groups associated to different choices are isomorphic. The group $H^1_G(\Gamma^\bullet, \underline{\mathbb{T}})$ classifies the G-equivariant complex line bundles on $\mathbb{T} \times M$. Then a G-equivariant complex line bundle λ_G is determined by a cocycle with components $h_{0,1}$ and $h_{1,0}$ in the double complex but the only nonidentity component is $h_{0,1} \in \prod_a C^\infty(N^{(1)}_a, \mathbb{T})$. Since $\delta(h_{0,1}) = 1$ we can consider $h_{0,1}$ as a sequence of functions $h_{ij} : \Gamma^{(1)} \to \mathbb{T}$. Then the cocycle relation can be written by $\partial^*(h) = 0$ which means

$$h_{ij}(g, hx) h_{jk}(h, x) = h_{ik}(gh, x).$$

The cocycle determines a G-equivariant complex line bundle by letting G act over each U_{ij} so that $(x_j, v) \mapsto ((gx)_i, h_{ij}(g, x)v)$ where (x_j, v) is written in the trivialization over U_j and $((gx)_i, h_{ij}(g, x)v)$ over U_i. The symmetric components h_{ii} define the group action on the fibres on

U_i. Associativity of the product is exactly the cocycle condition $\partial^*(h) = 0$.

3.4. Next we define the Fock bundle over $\mathbb{T} \times M$ in the spirit of [11] or 2.1. The twisting is now determined by the G-equivariant line bundle λ_G on M. We denote by $h \in H^1(\Gamma^\bullet, \underline{\mathbb{T}})$ a cohomology class of a G-equivariant line bundle λ_G constructed as in 3.3.

We define the operators S and N acting on the projective Fock space as in 2.1. Again, S is naturally defined up to fixing of a basis and therefore it is naturally defined as a $PU(\mathcal{H})$ operator. Then we define a projective Fock bundle associated to the cocycle $g \in H^1(\Gamma^\bullet, \underline{PU}(\mathcal{H}))$ with the nonidentity components defined by

$$\begin{aligned}
g_{ij}(g,x) &= (h_{ij}(g,x))^N, \\
g_{i+n,j+n}(g,x) &= (h_{i+n,j+n}(g,x))^N, \\
g^{(-1)}_{i,j+n}(g,x) &= (h_{i,j+n}(g,x))^N \\
g^{(-1)}_{j+n,i}(g,x) &= (h_{j+n,i}(g,x))^N \\
g^{(1)}_{i,j+n}(g,x) &= (h_{i,j+n}(g,x))^N S \\
g^{(1)}_{j+n,i}(g,x) &= S^{-1}(h_{j+n,i}(g,x))^N
\end{aligned}$$

for all $1 \leq i,j \leq n$. Here we denote $(g,x) \in G \times (\mathbb{T} \times M)$. The translation around the circle in the positive direction adds the charge by one and the translation in the negative direction subtract the charge by one. In local coordinates the action of G reads

$$(x_b, \Psi) \mapsto ((gx)_a, g_{ab}(g,x)\Psi) \qquad \text{for } 1 \leq a, b, \leq 2n$$

if (x_b, Ψ) is evaluated in the trivialization over U_b and $((gx)_a, \widehat{g}_{ba}(g,x)\Psi)$ in the trivialization over U_a and Ψ is a vector in the Fock space.

We denote by $\widehat{g}_{ab} \in H^1(\Gamma^\bullet, \underline{U}(\mathcal{H}))$ the lifted transition functions. The lifted transitions act on the Fock states over M so that each charge k subbundle restricted on M has a strcuture of a G-equivariant vector bundle of type $\lambda_G^{\otimes k}$. Over $\mathbb{T} \times M$ there is a G-equivariant gerbe. The Dixmier-Douady class $f \in H^2(\Gamma^\bullet, \underline{\mathbb{T}})$ is determined by the equations

$$f_{ijk}(g, hx, h, x) = \widehat{g}_{ij}(g, hx)\widehat{g}_{jk}(h,x)(\widehat{g}_{ik}(gh,x))^{-1}$$

for the lifted transition functions. Its nonidentity components are

$$\begin{aligned}
f^{(1)}_{i,j+n,k+n}(g,hx,h,x) &= (h_{j+n,k+n}(h,x))^{-1}, \\
f^{(1)}_{j+n,i,k+n}(g,hx,h,x) &= (h_{j+n,k+n}(gh,x)), \\
f^{(1)}_{j+n,k+n,i}(g,hx,h,x) &= (h_{j+n,k+n}(g,hx))^{-1},
\end{aligned} \tag{7}$$

for $1 \leq i,j,k \leq n$.

3.5. Let β_G denote the cocycle in $H^2(\Gamma^\bullet, \mathbb{Z})$ which maps to $h \in H^1(\Gamma^\bullet, \underline{\mathbb{T}})$ by the usual isomorphism and h determines the transtion functions of the G-equivariant bundle λ_G. Define $\alpha \in H^1(\Gamma^\bullet, \mathbb{Z})$ to be the groupoid cocycle with the nonzero components

$$\alpha^{(1)}_{i,j+n}(g,x) = 1 \qquad \text{and} \qquad \alpha^{(1)}_{i+n,j}(g,x) = -1$$

for all $1 \leq i,j \leq n$ and $(g,x) \in \Gamma$. At this level we can view α and β_G as groupoid de Rham cocycles. There is a cup product in $H^*_{\mathrm{dR}}(\Gamma^\bullet, \mathbb{Z})$, [2].

Proposition. The DD class $f \in H^2(\Gamma^\bullet, \mathbb{Z})$ is equivalent to the image of the cup product $\alpha \smile \beta_G$ under the usual group isomorphism $H^3(\Gamma^\bullet, \mathbb{Z}) \to H^2(\Gamma^\bullet, \underline{\mathbb{T}})$.

Proof. Denote by $h' \in H^1(\Gamma^\bullet, i\underline{\mathbb{R}})$ the cohomology class such that $\frac{1}{2\pi i}\partial^*(h') = \beta_G$ and $\exp(h') = h$. The nonidentity components of $\alpha \smile \beta_G$ are

$$(\alpha \smile \beta)^{(1)}_{i,j+n,k,l}(g_3, g_2g_1x, g_2, g_1x, g_1, x) = \frac{1}{2\pi i}(\partial^* h')^{(1)}_{j+n,k,l}(g_2, g_1x, g_1, x)$$

and

$$(\alpha \smile \beta)^{(1)}_{i+n,j,k,l}(g_3, g_2g_1x, g_2, g_1x, g_1, x) = -\frac{1}{2\pi i}(\partial^* h')^{(1)}_{jkl}(g_2, g_1x, g_1, x)$$

for $1 \leq i, j \leq n$ and $1 \leq k, l \leq 2n$. The cup product $\alpha \smile \beta_G$ is equal to $\frac{1}{2\pi i}(\partial^* t)$ where $t \in H^2(\Gamma^\bullet, i\underline{\mathbb{R}})$ is the cocycle with the nonzero components

$$t^{(1)}_{i,j+n,k}(g_2, g_1x, g_1, x) = -(h')^{(1)}_{jk}(g_1, x) \quad \text{and}$$
$$t^{(1)}_{i+n,j,k}(g_2, g_1x, g_1, x) = (h')^{(1)}_{jk}(g_1, x)$$

for $1 \leq i, j \leq n$ and $1 \leq k \leq 2n$. The image of the cup product in the standard isomorphism $H^3(\Gamma^\bullet, \mathbb{Z}) \to H^2(\Gamma^\bullet, \underline{\mathbb{T}})$ is the exponential cocycle of t.

We introduce another cochain, $d \in C^1(\Gamma^\bullet, \underline{\mathbb{T}})$ whose nonzero components are

$$d^{(1)}_{i+n,j}(g, x) = h_{i+n,j}(g, x)$$

for all $1 \leq i, j \leq n$. Then one checks that

$$f = \partial^*(d)\exp(t)$$

It follows that f is equal to $\alpha \smile \beta_G$ in cohomology. □

It follows that the DD-class is nontrivial since it can be realized as a cup product of two nontrivial cocycles which are localized on different manifolds. Moreover, we note that if we cut the circle $\mathbb{T}$, then the DD-class becomes trivial since the one cohomology class trivializes.

3.6. Consider the G-equivariant projective Fock bundle $\mathbf{PF}$ associated to the transition cocycle $g \in H^1(\Gamma^\bullet, \underline{PU}(\mathcal{H}))$ which is constructed as in 3.4. Again we fix a G-equivariant complex vector bundle ξ_G on M and then define the projective bundle $\mathbf{P}(\mathbf{F} \otimes \xi_G)$. We can assume that λ_G and ξ_G are trivialized over the U_i.

Following 2.2. we also define the spinor modules, $\mathcal{H}_s$. Then there are trivial Hilbert bundles $U_a \times (\mathcal{F} \otimes \mathcal{H}_s \otimes \mathbb{C}^r)$ where r is the rank of ξ_G. Then we define the local families $Q^a : U_a :\to \mathbf{Fred}^{(1)}_\Psi$ by

$$Q^a_{(\phi,p)} = (\sum_{n\in\mathbb{Z}} \psi_n \otimes e_n + \frac{\phi}{2\pi}\psi_0 \otimes \mathbf{1}) \otimes \mathbf{1},$$

The operators e_n are defined as in 2.1. It follows from the invariance of the cover that

$$g_{ab}(g, x)Q^b_{\phi,p}(g_{ab}(g^{-1}, gx))^{-1} = Q^a_{\phi-2\pi,gp} = Q^a_{g(\phi-2\pi,p)}.$$

for all $(\phi, p) = x \in U_{ab} \times M$ and $g \in G$. Therefore, Q is a G-invariant section of a $\mathbf{Fred}^{(1)}_\Psi$-bundle which is an associated bundle to $\mathbf{P}(\mathbf{F} \otimes \xi_G)$. Equivalently, Q is a section of a $\mathbf{Fred}^{(1)}_\Psi$-bundle over the action groupoid Γ and thus defines an element in the group $K^1(\Gamma, g)$.

Next we study the group $K^1(\Gamma, g)$. We use the closed cover $\mathbb{T} \times M = (\overline{\mathbb{T}}_+ \times M) \cup (\overline{\mathbb{T}}_- \times M)$ where $\overline{\mathbb{T}}_\pm$ is a closure of $\mathbb{T}_\pm$. Recall that the Dixmier-Douady class trvializes when we cut the circle $\mathbb{T}$. Thus, all the twisted K-theory groups of the components of the closed cover and their intersection are isomorphic to corresponding ordinary G-equivariant K-theory groups. Therefore, there is the following Mayer-Vietoris sequence associated to the action groupoid (see [20] 3.12.)

$$\begin{tikzcd}
K^0(\Gamma, g) \arrow[r, "c_0"] & K^0_G(\overline{\mathbb{T}}_+ \times M) \oplus K^0_G(\overline{\mathbb{T}}_- \times M) \arrow[r, "a_0"] & K^0_G(\overline{\mathbb{T}}_{+-} \times M) \arrow[d, "b_0"] \\
K^1_G(\overline{\mathbb{T}}_{+-} \times M) \arrow[u, "b_1"] & K^1_G(\overline{\mathbb{T}}_+ \times M) \oplus K^1_G(\overline{\mathbb{T}}_- \times M) \arrow[l, "a_1"'] & K^1(\Gamma, g) \arrow[l, "c_1"']
\end{tikzcd}$$

In twisted K-theory the maps a_* are defined by $a_*(x,y) = (x-y, x-y\otimes\lambda_G)$ where λ_G is the equivariant twisting line bundle.

The solution of $K^*(\Gamma, g)$ from this sequence is parallel to [11]. If λ_G is a nontorsion element in $H^2_G(M)$, then there is a group isomorphism

$$K^*(\Gamma, g) \simeq K^{*-1}_G(M)/(K^{*-1}_G(M)\otimes(1-\lambda_G))$$

In the general case when λ_G is nontrivial $K^*(\Gamma, g)$ is isomorphic to an extension of the group

$$\{x\in K^*_G(M) : x = x\otimes\lambda_G\} \quad \text{by} \quad K^{*-1}_G(M)/(K^{*-1}_G(M)\otimes(1-\lambda_G)).$$

In the latter case, one cannot fix the group extension from the Mayer-Vietoris sequence.

3.7. The hyper cohomology groups $H^k(\Gamma^\bullet, \mathbb{Z})$ are isomorphic to the G-equivariant integer cohomology groups because we are working with a groupoid Morita equivalent to an action groupoid. We shall use the Cartan model of differential equivariant cohomology to study equivariant superconnections associated to the K^1-group elements of 3.6.

Let $\mathfrak{g}$ denote the Lie algebra of G. If $X\in\mathfrak{g}$ we denote by X_M the vector field induced by the action $\exp(-tX)$ on M. Denote by $\iota(X_M) : \Lambda^\bullet(M)\to\Lambda^{\bullet-1}(M)$ the contraction by X_M. Define

$$\Lambda^q_G(M) = \bigoplus_{2i+j=q} (S^i(\mathfrak{g}^*)\otimes\Lambda^j(M))^G,$$

the order q subspace in the space of equivariant polynomial functions $\mathfrak{g}\to\Lambda^*(M)$. The order zero subspace consists of G-equivariant functions. The equivariant coboundary $d_G : \Lambda^q_G(M)\to\Lambda^{q+1}_G(M)$ is defined by

$$(d_G\alpha)(X) = d(\alpha(X)) - \iota(X_M)(\alpha(X))$$

The square $d^2_G\alpha(X) = -\{d, \iota(X_M)\}\cdot\alpha(X)$ is equal to zero since $\alpha(X)$ is invariant under the action of the one parameter group generated by X. We denote by $H^*_G(M, \mathbb{R})$ the cohomology of this complex.

If ∇ is an invariant connection of an equivariant line bundle λ_G, then the equivariant curvature of ∇ is $F(X) = (\nabla - \iota(X_M))^2 + \mathcal{L}(X)$ where $\mathcal{L}(X)$ is the Lie derivative of the action of G, see [4], [10]. The first Chern class, c^G_1 getting values in $H^2_G(M, \mathbb{Z})$ sends an isomorphism class of a G-invariant line bundle to the cohomology class of its equivariant curvature form which is an integer class. The tensor product of equivariant line bundles defines a group structure in the set of isomorphism classes of G-equivariant line bundles. Then c^G_1 an isomorphism of groups.

Let ∇_ξ and ∇_λ be G-invariant connections of the bundles ξ_G and λ_G. We pullback the projective bundle $\mathbf{P}(\mathbf{F}\otimes\xi_G)$ to the covering space $\mathbb{R}\times M$. Moreover, we pullback the twisted K^1-theory elements Q to $\mathbb{R}\times M$. The DD class becomes trivial and therefore, we can lift its transition functions to unitary operators. Let us denote by $\mathbf{F}_\xi$ the Hilbert bundle on $\mathbb{R}\times M$. Then a connection of $\mathbf{F}_\xi$ can be chosen by $\nabla = e_0\nabla_\lambda + \nabla_\xi$. $\hat{F}$ denotes its curvature. Notice that e_0, defined in 2.2., counts the charge of the Fock state it acts. We also consider the homotopy $t\mapsto\sqrt{t}Q = Q_t$ for the parameter $t\geq 0$. We define an equivariant superconnection by $D_t = \chi Q_t + \nabla$. The equivariant super curvature is defined by

$$\mathbb{F}_t(X) = Q^2_t - \chi[\nabla, Q_t] + \hat{F}(X)$$

for all $X\in\mathfrak{g}$. We have introduced a formal symbol χ satisfying $\chi^2 = 1$ which anticommutes $\iota(X_M)$ and equivariant forms of odd rank and commutes with Q. We define an equivariant super Chern character by

$$\Theta_t(X) = \mathrm{sTr}(\exp(-\mathbb{F}_t(X))).$$

sTr is the supertrace which applies the usual operator trace to the terms linear in χ and maps to zero those which are not.

Proposition. The following holds for the equivariant superconnection on $\mathbb{R}\times M$.

(1) The form is closed in the equivariant cohomology: $d_G\Theta_t(X) = 0$.

(2) Its cohomology class is independent of t.

Proof. (1) If $\phi \in \Lambda^*_G(\mathbb{T} \times M) \otimes \mathrm{End}(\mathbf{F}_\xi)$ is such that $d_G\phi = (d_G \cdot \phi) + \phi d_G$, then we have $d_G(\mathrm{sTr}(\phi)) = \mathrm{sTr}[d_G, \phi]$. In local coordinates we see that

$$\mathbb{F}_t(X) = Q_t^2 - \chi[d + \omega, Q_t] + \hat{F}(X) = Q_t^2 - \chi(d \cdot Q_t) + \hat{F}(X).$$

Then

$$\begin{aligned}
[d_G, \mathbb{F}_t(X)] &= [d_G, Q_t^2 - \chi(d \cdot Q_t)] \\
&= [d - \iota(X_M), Q_t^2] - \chi\{d - \iota(X_M), (d \cdot Q_t)\} \\
&= (d \cdot Q_t^2) + \chi(\iota(X_M)d) \cdot Q_t \\
&= (d \cdot Q_t^2) - \chi(d\iota(X_M)) \cdot Q_t \\
&= (d \cdot Q_t^2) = 2(d \cdot Q_t)
\end{aligned}$$

where we have used that Q_t is an invariant form and therefore $\{d_G, \iota(X_M)\} \cdot Q_t = 0$. It follows

$$\begin{aligned}
\mathrm{sTr}[d_G, (\mathbb{F}_t(X))^n] &= \mathrm{sTr}(n[d_G, \mathbb{F}_t(X)]\mathbb{F}_t(X)) \\
&= \mathrm{sTr}(2n(d \cdot Q_t)\mathbb{F}_t(X)^{n-1})
\end{aligned}$$

Recall that the supertrace only picks components linear in χ. If $n \geq 2$ in the above formula, one can use the commutation relations to verify that supertrace only picks terms which contain $(d \cdot Q_t)^k$ for some $k \geq 2$. Thus supertrace is identically zero. In the case $n = 1$ supertrace is zero because there are no terms linear in χ.

(2) Following the lines of [19] it is straightforward to check that the following transgression formula holds

$$\frac{d}{dt}\Theta_t(X) = d_G \mathrm{sTr}(\frac{dQ_t}{dt}\Theta_t(X)). \qquad \square$$

We will need the following.

Proposition. $[F_\lambda(X), \mathbb{F}_t(X)] = 0$ for all $X \in \mathfrak{g}$ and $t \in \mathbb{R}$.

Proof. The equivariant curvature forms commute with each other, thus $[F_\lambda(X), \hat{F}(X)] = 0$. Then we apply the local coordinates to obtain

$$\begin{aligned}
[F_\lambda(X), Q] &= [\nabla_\lambda^2 - \{\nabla_\lambda, \iota(X)\} + \mathcal{L}(X), Q_t] \\
&= [-\{d + \omega_\lambda, \iota(X)\}, Q_t] \\
&= [-\{d, \iota(X)\}, Q_t] - [\iota(X)(\omega_\lambda), Q_t] = 0.
\end{aligned}$$

where we have used that Q_t is G-invariant and therefore the Lie algebra action commutes with Q_t and $\iota(X)$ and d are antiderivations and $\{d_G, \iota(X)\} \cdot Q_t = 0$. Similarly,

$$\begin{aligned}
[F_\lambda(X), \chi(d \cdot Q_t)] &= \chi[\nabla_\lambda^2, d \cdot Q_t] - \chi[\{d + \omega_\lambda, \iota(X)\}, d \cdot Q_t] + \chi[\mathcal{L}(X), d \cdot Q_t] \\
&= \chi[\{d, \iota(X)\}, (d \cdot Q_t)] = 0
\end{aligned}$$

because Q_t is G invariant and so is $d \cdot Q_t = (d + \iota(X)) \cdot Q_t$ and therefore also in the kernel of $\{d, \iota(X)\}$. $\square$

Consider $\hat{F}(X)$ as a G-equivariant connection of $\mathbf{F}_\xi$ on $\mathbb{R} \times M$. The connection depends on the parameter $y \in \mathbb{R}$ since the charge grows under the translations by 2π. Therefore, we cannot push this form to the space $\Lambda_G^{\mathrm{odd}}(\mathbb{T} \times M)$ since it would depend on the choice of the angle variable. However, since the equivariant curvature $F_\lambda(X)$ commutes with $\Theta_t(X)$, we push the form to the quotient space

$$\Lambda_G^{\mathrm{odd}}(\mathbb{T} \times M)/(\Theta_t \wedge F_\lambda).$$

Putting everything together we see that for each $t \in \mathbb{R}_+$ the equivariant super Chern character defines a map

$$\Theta_t : K^1(\Gamma, g) \to H_G^{\mathrm{odd}}(\mathbb{T} \times M)/(\Theta_t \wedge F_\lambda).$$

We use this map to distinguish different group elements in the equivariant twisted K-theory group. As usual, we study the limit $t \to \infty$ to simplify the cohomology classes. We choose the superconnection with the following curvature

$$\mathbb{F}_t(X) = tQ^2 - \sqrt{t}\chi(d \cdot Q) + \hat{F}(X) = tQ^2 - \sqrt{t}\chi\psi_0 dy + \hat{F}(X)$$

We can proceed as in [11]. The limit is equal to

$$\lim_{t \to \infty} \Theta_y = \sqrt{\pi}\delta(e_0 + y)\mathrm{tr}_\xi(e^{-\hat{F}(X)}) \wedge dy,$$

where δ is the Dirac delta distribution. This form can be pushed down to the quotient space cohomology group.

4 Examples

4.1. In the first part of this series, [11], we gave explicit construction for the twisted K^1-groups on the product manifolds $\mathbb{T} \times M$ when $M = \mathbb{T}^2$ and S^2. In both cases, the usual K-theory groups can be computed, for example, using the Kunneth formula which leads to

$$K^1(\mathbb{T} \times S^2) \simeq \mathbb{Z}^{\oplus 2}, \qquad K^1(\mathbb{T} \times \mathbb{T}^2) \simeq \mathbb{Z}^{\oplus 4}.$$

The twisting line bundles are classified by integers. For λ corresponding to $0 \neq k \in \mathbb{Z}$ and ξ to $0 \neq j \in \mathbb{Z}$ the twisted K^1-groups are

$$K^1(\mathbb{T} \times M) \simeq \mathbb{Z} \oplus \mathbb{Z}_k.$$

The torsion sugroups appear in the twisted theory. In the following we study a phenomenon contrary to this one.

4.2. We study the real projective space $\mathbb{R}P^{2n}$ for $n \in \mathbb{N}$. No group action on $\mathbb{R}P^{2n}$ is assumed here. The second cohomology group is the torsion group $\mathbb{Z}_2$. Let us denote by λ the nontrivial line bundle which is subject to the relation $\lambda \otimes \lambda = 1$. The cohomology groups of $\mathbb{T} \times \mathbb{R}P^{2n}$ can by computed with Kunneth's formula. The third cohomology is

$$H^3(\mathbb{T} \times \mathbb{R}P^{2n}) = \mathbb{Z}_2 \quad \text{for all } n \in \mathbb{N}.$$

There is exactly one nontrivial twisting class $\sigma = \alpha \smile \beta$ where β is the class of λ in $H^2(\mathbb{R}P^2)$ and α is the generator of $H^1(\mathbb{T})$.

K-theory groups on $\mathbb{R}P^{2n}$ are solved in [1]

$$K^0(\mathbb{R}P^{2n}) \simeq \mathbb{Z} \oplus \mathbb{Z}_{2^n}, \qquad K^1(\mathbb{R}P^{2n}) = 0.$$

As an abelian group, K^0 is generated by 1 and x_n and its ring structure is given by

$$x_n = 1 - [\lambda], \qquad x_n^2 = 2x_n, \qquad 2^n x_n = 0. \tag{8}$$

By Kunneth's formula we have

$$K^1(\mathbb{T} \times \mathbb{R}P^{2n}) \simeq \mathbb{Z} \oplus \mathbb{Z}_{2^n}.$$

Next we use the results of 3.6. to solve the twisted K^1-group. The normal subgroup $K^0(\mathbb{R}P^{2n}) \otimes (1 - \lambda) = K^0(\mathbb{R}P^{2n}) \otimes x_n$ is the torsion part $\mathbb{Z}_{2^n}$ of $K^0(\mathbb{R}P^{2n})$ by (8). Therefore, the σ-twisted K^1-group is given by

$$K^1(\mathbb{T} \times \mathbb{R}P^{2n}, \sigma) = \mathbb{Z}.$$

We observe that the K^1-theory group of $\mathbb{T} \times \mathbb{R}P^{2n}$ has a torsion subgroup for any $n \in \mathbb{N}$ and the twisting by σ kills the torsion subgroup.

Consider a pullback of the projective bundle of Fock spaces $\mathbf{PF}_\xi$ to $\mathbb{R} \times \mathbb{R}P^{2n}$ as in 3.7. Then we can lift the transition functions and define a Hilbert bundle on the covering space. Assume that ξ is a line bundle on $\mathbb{R}P^{2n}$. Thus, $\xi = \lambda$ or $\xi = 1$, the trivial complex line bundle. Then the topological type of the vacuum at the submanifold with $0 = y \in \mathbb{R}$ is ether λ or 1. The type depends on the chosen lift $\mathbb{R} \to \mathbb{T}$ but is not important. Translations in the direction of positive real numbers by one then change this type to 1 or λ (respectively). This continues through the real line: the topological type of the vacuum is $\mathbb{Z}_2$ graded and depends on the choice of integer part of the $\mathbb{R}$ coordinate. This is the reason for the absence of the torsion subgroup: twisted K^1-theory does not distinguish λ from 1 since vacuums of these types are related by a choice of the phase of the angle variable over $\mathbb{T}$ which is arbitrary.

Going back to the nontwisted case (i.e. set $\beta = 0$), we have the Fredholm operator construction on $\mathbb{R} \times \mathbb{R}P^{2n}$ and the kernels are localized over integers but the topological type of this bundle is constant in $\mathbb{R}$. Therefore, considered as a family over $\mathbb{T} \times \mathbb{R}P^{2n}$, the topological type of a vacuum line bundle does matter.

Now suppose that ξ is an arbitrary complex vector bundle. We can study the super Chern character associated to the local families Q, as in 3.7. with a trivial group action G. Detailed calculations can also be found from Section 7 in [11]. The curvature form of λ vanishes in the de Rham cohomology. Then the Chern character gets values in the usual odd de Rham cohomology of $\mathbb{T} \times \mathbb{R}P^{2n}$. In fact, according to the analysis in 3.7. the image of the Chern character in the limit $t \to \infty$ is equal to the push forward of $\sqrt{\pi}\delta(e_0 + y)\mathrm{rk}(\xi)$ on $\mathbb{T} \times \mathbb{R}P^{2n}$ and therefore the rank of the vacuum bundle determines the twisted K^1-class. We conclude that the super Chern character is a noninjective group homomorphism $K^1(\mathbb{T} \times \mathbb{R}P^{2n}) \to H^{\mathrm{odd}}(\mathbb{T} \times \mathbb{R}P^{2n})$ but it is a group isomorphism $K^1(\mathbb{T} \times \mathbb{R}P^{2n}, \sigma) \to H^{\mathrm{odd}}(\mathbb{T} \times \mathbb{R}P^{2n})$ for $\sigma = \alpha \smile \beta$.

4.3. Let $\mathbb{T}$ act on S^2 by rotations around the axis of the sphere. We denote by S^2_+ and S^2_- the north and south hemispheres of S^2 such that the intersection S^2_{+-} is the equator $\mathbb{T}$. Then $S^2 = S^2_+ \cup S^2_-$ and the cover is $\mathbb{T}$-invariant. The south and north poles are the fixed points of this action. The equivariant line bundles are classified by

$$H^2_{\mathbb{T}}(S^2, \mathbb{Z}) \simeq \mathbb{Z}[u_+] \oplus \mathbb{Z}[u_-].$$

This result is well known and can be solved with the Mayer-Vietoris sequence. $u_\pm$ are the generators of the cohomology groups $H^2_{\mathbb{T}}(S^2_\pm, \mathbb{Z})$: $S^2_\pm$ are retracted to the fixed point and these groups are isomorphic to the $\mathbb{T}$-equivariant cohomology of a point, i.e. $H^2(\mathbb{C}P^\infty, \mathbb{Z})$ and so under this identification $u_\pm$ can be see as generators of $H^2(S^2, \mathbb{Z})$. We also have

$$H^3_{\mathbb{T}}(\mathbb{T} \times S^2, \mathbb{Z}) \simeq H^2_{\mathbb{T}}(S^2, \mathbb{Z})$$

by Mayer-Vietoris sequence.

Suppose that $\lambda_{\mathbb{T}}$ is a $\mathbb{T}$-equivariant complex line bundle over M associated to the class (n_+, n_-) in $H^2_{\mathbb{T}}(S^2, \mathbb{Z})$. Then $\mathbb{T}$ acts on the fibers over $S^2_\pm$ through the character $\exp(i\varphi) \mapsto \exp(in_\pm\varphi)$.

The ordinary equivariant K-theory can be computed using the Mayer-Vietoris sequence

$$\begin{tikzcd}
K^0_{\mathbb{T}}(S^2) \arrow[r, "c_0"] & K^0_{\mathbb{T}}(S^2_+) \oplus K^0_{\mathbb{T}}(S^2_-) \arrow[r, "a_0"] & K^0_{\mathbb{T}}(S^2_{+-}) \arrow[d, "b_0"'] \\
K^1_{\mathbb{T}}(S^2_{+-}) \arrow[u, "b_1"] & K^1_{\mathbb{T}}(S^2_+) \oplus K^1_{\mathbb{T}}(S^2_-) \arrow[l, "a_1"'] & K^1_{\mathbb{T}}(S^2) \arrow[l, "c_1"']
\end{tikzcd}$$

The groups $K^0_{\mathbb{T}}(S^2_+)$ and $K^0_{\mathbb{T}}(S^2_-)$ are isomorphic to $K^0_{\mathbb{T}}(*)$ where $*$ is a pole and $K^0_{\mathbb{T}}(*)$ is isomorphic to the representation ring

$$R(\mathbb{T}) = \mathbb{Z}[a, a^{-1}].$$

The circle group action on S^2_{+-} is free. Therefore

$$K^0_{\mathbb{T}}(S^2_{+-}) \simeq K^0(\mathbb{T}/\mathbb{T}) \simeq \mathbb{Z} \qquad \text{and}$$
$$K^1_{\mathbb{T}}(S^2_{+-}) \simeq K^1(\mathbb{T}/\mathbb{T}) \simeq 0$$

Next we observe that $K^1_{\mathbb{T}}(S^2_+) \simeq K^1_{\mathbb{T}}(*)$. Thus, $K^1_{\mathbb{T}}(S^2_+) = 0$ and similarly $K^1_{\mathbb{T}}(S^2_-) = 0$. The homomorphisms a_i, $i = 0, 1$ sends (x, y) to $x - y$. The group elements in $K^0_{\mathbb{T}}(S^2_{+-})$ are determined by an integer, the virtual rank. Thus,

$$\mathrm{Ker}(a_0) \simeq (R(\mathbb{T}) \oplus R(\mathbb{T}))/\mathbb{Z}, \qquad \mathrm{Im}(a_0) \simeq \mathbb{Z} \quad \text{and} \quad \mathrm{Ker}(a_1) = 0.$$

Since there is no torsion in these groups the K-theory groups are as follows

$$\begin{aligned} K^0_{\mathbb{T}}(S^2) &\simeq (K^1_{\mathbb{T}}(S^2_{+-})/\mathrm{Im}(a_1)) \oplus \mathrm{Ker}(a_0) \simeq (R(\mathbb{T}) \oplus R(\mathbb{T}))/\mathbb{Z} \\ K^1_{\mathbb{T}}(S^2) &\simeq (K^0_{\mathbb{T}}(S^2_{+-})/\mathrm{Im}(a_0)) \oplus \mathrm{Ker}(a_1) \simeq 0. \end{aligned}$$

The elements in $K^0_{\mathbb{T}}(S^2)$ are the pairs $(x, y) \in R(\mathbb{T})^{\oplus 2}$ constrained by the common virtual dimension. Topologically any vector bundle over S^2 is a direct summand of line bundles. Therefore, $K^0_{\mathbb{T}}(S^2)$ is the abelian group generated by the $\mathbb{T}$-equivariant complex line bundles and each generator carries a representations $(a^{n_+}, a^{n_-}) \in R(\mathbb{T})^{\oplus 2}$ on its fibres.

The twisted K^1-group can be computed from the formulas in 3.6. If $\lambda_{\mathbb{T}} = (l_+, l_-)$ is an equivariant line bundle and the K^1-group is subject to the twisting of type $\alpha \smile \beta_{\mathbb{T}}$ and $\beta_{\mathbb{T}}$ is the 2-cohomology class which identifies with the equivalence class of $\lambda_{\mathbb{T}}$, then we have

$$K^1_{\mathbb{T}}(S^2, \widehat{g}) = K^0_{\mathbb{T}}(S^2)/K^0_{\mathbb{T}}(S^2) \otimes (1 - \lambda_{\mathbb{T}}).$$

Therefore, we observe that the twisted K-group is the abelian group defined by

$$K^1_{\mathbb{T}}(S^2, g) = K^0_{\mathbb{T}}(S^2)/\{(a^{n_+}, a^{n_-}) \sim (a^{n_+ + l_+}, a^{n_- + l_-})\}.$$

We can find explicit groupoid cocycles which determine the $\mathbb{T}$-equivariant gerbes and twisted K^1-classes. Now we replace the closed sets $S^2_\pm$ with $\mathbb{T}$-invariant open sets which cover the north and the south hemispheres and which intersect in a tube around the equator. These sets are still denoted by $S^2_\pm$. The equivariant complex line bundle corresponding to (a^{n_+}, a^{n_-}) in $H^2_{\mathbb{T}}(S^2)$ is associated to the groupoid cocycle

$$h \in H^1((\coprod_{ij} \mathbb{T} \times S^2_{ij})^\bullet, \underline{\mathbb{T}})$$

for $i, j = +, -$ and its components are

$$\begin{aligned} &h_{++}(\exp(i\varphi), p) = \exp(in_+\varphi), \\ &h_{--}(\exp(i\varphi), p) = \exp(in_-\varphi), \\ &h_{+-}(\exp(i\varphi), (\alpha, r)) = \exp(in_+\varphi + i(n_+ - n_-)\alpha) \qquad \text{and} \\ &h_{-+}(\exp(i\varphi), (\alpha, r)) = \exp(in_-\varphi - i(n_+ - n_-)\alpha) \end{aligned}$$

where we have denoted by α the angle coordinate and r the height coordinate of the tube around the equator S^2_{+-}. The components h_{++} and h_{--} carry the information of the $\mathbb{T}$-action on the fibres. The topological type of this bundle is determined by $h_{+-}(1, \alpha)$ which is a function on the equator with winding number equal to $(n_+ - n_-)$. The cocycle g of the projective Fock bundle is straightforward to write down using 3.4.

Next we set $\xi_{\mathbb{T}}$ to be the line bundle corresponding to the class (a^{j_+}, a^{j_-}). Then we apply the superconnection analysis on the covering space $\mathbb{R} \times S^2$ to study the dependence of the choice of $\xi_{\mathbb{T}}$ in twisted K-theory. We have

$$\lim_{t \to \infty} \Theta_t(X) = \sqrt{\pi}\delta(e_0 + y)dy \wedge (1 - \hat{F}(X)).$$

Next we push this form to $\mathbb{T} \times M$. Let F_ξ and F_λ denote the G-invariant connections. Then the three form component is cohomologous to

$$\begin{aligned} &-\frac{\sqrt{\pi}dy \wedge F_\xi}{2\pi} \quad \mathrm{mod} \quad \frac{\sqrt{\pi}dy \wedge F_\lambda}{2\pi} \quad \Leftrightarrow \\ &-\frac{\sqrt{\pi}dy}{2\pi} \wedge (j^+F^+ + j^-F^-) \quad \mathrm{mod} \quad \frac{\sqrt{\pi}dy}{2\pi} \wedge (l^+F^+ + l^-F^-) \end{aligned}$$

The equivariant forms $F^{\pm}$ are generators of $H^2_{\mathbb{T}}(S^2,\mathbb{Z})$. Therefore, if ξ_1 and ξ_2 are equivariant line bundles associated to the classes $(a^{j_1^+},a^{j_1^-})$ and $(a^{j_2^+},a^{j_2^-})$, and $\lambda_{\mathbb{T}}$ is as above, then the families Q acting on $\mathbf{F}_{\xi_1}$ and $\mathbf{F}_{\xi_1}$ are different elements of $K^1_{\mathbb{T}}(\mathbb{T}\times S^2,g)$ if and only if

$$j_1^+ - j_2^+ \neq 0 \quad \mathrm{mod} \quad l^+ \quad \text{or} \quad j_1^- - j_2^- \neq 0 \quad \mathrm{mod} \quad l^-.$$

It is impossible to distinguish all the different twited K^1-theory classes in equivariant cohomology. If we apply similar analysis as above for higher rank $\mathbb{T}$-equivariant vector budles ξ_1 and ξ_2, the Chern character map is noninjective. Nevertheless, if the ranks of ξ_1 and ξ_2 do not match, then the associated K^1-classes are necessarily different since the one form components of the Chern characters are not equal in cohomology.

Appendix

A.1. This appendix is motivated by the geometric and group theoretic structures in Hamiltonian quantization. In quantum field theory on a unit circle $\mathbb{T}$, one defines a space of potentials $\mathcal{A}$ to be the affine space of $\mathfrak{g}$-valued one forms on $\mathbb{T}$ and $\mathfrak{g}$ is a Lie algebra of a simple, compact and simply connected Lie group G. Then there is a family of Dirac operators acting on $L^2(\mathbb{T})$ which are coupled to the potentials. The gauge symmetry group is chosen to be the group of based gauge transformations ΩG which is the subgroup in the group of smooth loops LG based at the unit $\gamma(0)=1=\gamma(2\pi)$. The true parameter space of the theory is the quotient $\mathcal{A}/\Omega G$ where ΩG acts on $\mathcal{A}$ by

$$(g,A)\mapsto gAg^{-1}+gd(g^{-1}).$$

The action makes $\mathcal{A}\to\mathcal{A}/\Omega G$ a principal bundle with a projection map sending $A\in\mathcal{A}$ to a holonomy around the circle.

On $\mathcal{A}/\Omega G$ which is homeomorphic to G we define a Fock bundle. We set vacuum levels x_i and a finite cover U_i of $\mathcal{A}/\Omega G$ so that the the numbers x_i do not belong to the spectrum of the Dirac family over U_i. Then, over each U_i one can define a polarization of the Hilbert space into Dirac eigenspaces with eigenvalues larger than x_i and its orthogonal complement. There is spectral flow and the vacuum cannot be fixed globally. On the overlaps U_{ij} two different vacuums associated to different polarizations are related by tensoring with a determinant line bundle. Then the Fock bundle is naturally a projective bundle of Fock spaces. Then there is a prolongation problem: can we fix the vacuum levels (determinant line bundles) such that the Fock bundle becomes a bundle of Hilbert spaces over $\mathcal{A}/\Omega G$. In [5], [13] it was found that the obstruction to do this is the three cohomology part of the odd Chern character of the Dirac family over $\mathcal{A}/\Omega G$. The twisted K-theory analysis in the first part of this series [11] was based on this result.

The gauge group ΩG acts on the fibers of the Fock bundle as a subgroup of LG. The loop group acts by an irredcible positive energy representation which is projective. Alternatively, these representations are irreducible representations of the central extension $\hat{LG}$. The extension cocycles arise in the lifted transition functions of the Fock bundle over $\mathcal{A}/\Omega G$ and ruin the Cech cocycle conditions. The relationship to the prolongation problem can be described explicitly. Let $c\in H^2(l\mathfrak{g},\mathbb{C})$ denote the Lie algebra cohomology class of the extension. We can identify the Lie algebra elements with the tangent vector fields near the identity and realize c as a differential two form near the identity of LG and extend it by left translation through LG. The normalized cocycle $c/2\pi i$ defines an integral cohomology class. We choose local sections $s_i:U_i\to p^{-1}(U_i)$ of $\mathcal{A}\xrightarrow{p}\mathcal{A}/\Omega G$. Let g_{ij} denote the ΩG valued transition functions of the principal bundle $\mathcal{A}$. Then $s_j=s_ig_{ij}$. We pull back the two forms c on LG to define local 2-forms $g_{ij}^*(c/2\pi i)$ on U_{ij}. These define the DD class of the gerbe in Cech-de Rham double complex. We find the forms ω_i and ω_j over U_i and U_j so that $g^*_{ij}(c/2\pi i)=\omega_i-\omega_j$. Let Ω be a 3-form over G which is locally defined by $d\omega_i$ on U_i. Then Ω is a global 3-form which represents the DD-class of the gerbe. Explicitly,

$$\Omega = k\frac{1}{24\pi^2}(g^{-1}dg)^3$$

is k times the generator of $H^3(G,\mathbb{Z})$ if the projective representation of LG has level equal to k, [16], [17].

A.2. We can view the torus $\mathbb{T}^3$ as a parameter space for a quantum field theory over $\mathbb{T}_\theta$: we replace $\mathcal{A}$ by $\mathbb{R}^3$ and ΩG by $\mathbb{Z}^3$. The action is by translations and therefore, $\mathbb{R}^3/\mathbb{Z}^3 = \mathbb{T}^3$. We write $\mathbb{T}^3 = \mathbb{T}_\phi \times \mathbb{T}^2$. In [11] we constructed a Hilbert bundle over the 4-torus $\mathbb{T}_\theta \times \mathbb{T}^3$ with fibers $L^2(\mathbb{T}_\theta)$. There is a Dirac family over $\mathbb{T}_\theta \times \mathbb{T}^3$ coupled to the connection of the bundle $\lambda_1 \boxtimes \lambda_k$ where λ_i is the line bundle associated to the Chern class i on $\mathbb{T}^2$. The 3 cohomology component of the Chern character of this Dirac family is the cup product $\alpha \smile \beta$ such that α is the generator of $H^1_{\mathrm{dR}}(\mathbb{T}_\phi, \mathbb{Z})$ and β is $(2\pi i)^{-1}$ times the curvature form of λ_k. The associated decomposable Cech cocycle defines a transition data of the gerbe exactly as in 2.1.

Here we want to connect the topological prolongation problem to a group extension problem: we can define a trivial bundle of Fock spaces on the affine space $\mathbb{R}^3$ and then implement an extended $\mathbb{Z}^3$-symmetry on the fibres by an extension cocycle which defines the cup product class in cohomology. In the following we apply the theory of groupoid extensions to make the connection between this extension problem and the cohomology class of the gerbe explicit.

A.3. We can consider the torus $\mathbb{T}^3$ as a transformation groupoid $\Gamma := \mathbb{Z}^3 \times \mathbb{R}^3 \rightrightarrows \mathbb{R}^3$ with the usual target and source maps

$$t, s : \Gamma \rightrightarrows \mathbb{R}^3, \qquad s(a,x) = x, \qquad t(a,x) = x + a.$$

The composition $m : \Gamma^{(2)} \to \Gamma$ is defined in its domain by $(b, x+a)(a,x) = (a+b, x)$. The action is free and therefore the cohomology of this groupoid is just the usual de Rham cohomology of the 3-torus.

Next we define a topologically trivial $\mathbb{T}$-extension $\hat{\Gamma} := \mathbb{Z}^3 \times \mathbb{R}^3 \times \mathbb{T} \rightrightarrows \mathbb{R}^3$ with $p : \hat{\Gamma} \to \Gamma$ defined by $p(a,x,\mu) = (a,x)$. The target and source maps of the groupoid $\hat{\Gamma}$ are

$$\widehat{s} = s \circ p, \qquad \widehat{t} = t \circ p.$$

The nontriviality of the extension is associated to the cocycle $\omega \in H^2(\Gamma^{(2)}, \underline{\mathbb{T}})$. The cocycle condition is

$$(\partial^*(\omega))((c, x+a+b), (b, x+a), (a,x)) = 0$$

which is equivalent to

$$\begin{aligned}&\omega((b, x+a), (a,x))\omega((c, x+a+b), (a+b, x)) = \\ &\omega((b+c, x+a), (a,x))\omega((c, x+a+b), (b, x+a)).\end{aligned}$$

We set $x = (x_0, x_1, x_2)$ and $a = (a_0, a_1, a_2)$. One checks easily that $\omega : \Gamma^{(2)} \to \mathbb{T}$ defined by

$$\omega((b, x+a), (a,x)) = \exp(2\pi i k a_0 b_2 x_1). \tag{9}$$

is a cocycle. Then we define a composition in the extension groupoid $\hat{\Gamma}$ by

$$(b, x+a, \mu')(a, x, \mu) = (a+b, x, \mu\mu'\omega((b, x+a), (a,x))).$$

This extensions corresponds to the class $k \in \mathbb{Z} \simeq H^3(\Gamma^\bullet, \mathbb{Z})$. This will be justified below.

There are the generators $\widetilde{\alpha}_i$ with $i = 0, 1, 2$ in $H^1(\Gamma^\bullet, \mathbb{Z}) \simeq \mathbb{Z}^3$ which are given by

$$\widetilde{\alpha}_i(a,x) = a_i \quad \text{for all} \quad (a,x) \in \mathbb{Z}^3 \times \mathbb{R}^3.$$

The cohomology classes of a 3-torus can be all constructed with the cup product. The generator of the 3-cohomology is the class

$$(\widetilde{\alpha}_2 \smile \widetilde{\alpha}_1 \smile \widetilde{\alpha}_0)((c, x+a+b), (b, x+a), (a,x)) = a_0 b_1 c_2.$$

If we choose $\widetilde{\beta}$ to be the class $k\widetilde{\alpha}_1 \smile \widetilde{\alpha}_2$, then we can write, by antisymmetry of the generators

$$(\widetilde{\alpha}_0 \smile \widetilde{\beta})((c, x+a+b), (b, x+a), (a,x)) = -k a_1 b_0 c_2.$$

Next we define a cocycle $f' \in H^2(\Gamma^\bullet, \underline{i\mathbb{R}})$ by

$$f'((b, x+a), (a, x)) = 2\pi i k a_0 b_2 x_1.$$

This satisfies $\frac{1}{2\pi i}\delta(f') = \widetilde{\alpha} \smile \widetilde{\beta}$. It then follows that under the usual isomorphism, $\widetilde{\alpha} \smile \widetilde{\beta}$ maps to the cocycle $\omega \in H^2(\Gamma^\bullet, \underline{\mathbb{T}})$ defined in (9).

In the action groupoid formalism there is a globally trivial $\mathbb{Z}^3$-equivariant projective vector bundle over $\mathbb{R}^3$ with a $H^1(\Gamma^\bullet, \underline{PU}(\mathcal{H}))$ class uniquely determined (in cohomology) by ω in (9). The equivariant bundle is topologically $\mathbb{R}^3 \times \mathcal{PF}$ where $\mathcal{PF}$ is the projective Fock space and the $\mathbb{Z}^3$ action is defined by

$$a \triangleright (x, \Psi) = (x + a, g(a, x)\Psi)$$

for some cocycle $g \in H^1(\Gamma^\bullet, \underline{PU}(\mathcal{H}))$. The cocycle condition $\partial^*(g) = 0$ is equivalent to the associativity of the action. If we lift the action to the group $U(\mathcal{H})$ then $\omega \in H^2(\Gamma^\bullet, \underline{\mathbb{T}})$ is risen from the relation

$$\partial^*(\widehat{g}) = \omega \qquad \Leftrightarrow \qquad \widehat{g}(b, x+a)\widehat{g}(a, x)\widehat{g}(a+b, x)^{-1} = \omega((b, x+a), (a, x))$$

where $\widehat{g}$ are the lifted functions $\Gamma^{(1)} \to U(\mathcal{H})$.

We define the cocycle g by (the operators S and N are defined as in 2.1.)

$$g(a, x) = S^{-a_0} \exp(-2\pi i k a_2 x_1)^N. \tag{10}$$

It is straightforward to check that g defines a cohomology class in $H^1(\Gamma^\bullet, \underline{PU}(\mathcal{H}))$. The DD class of g is given by

$$\omega((b, x+a), (a, x)) = \exp(2\pi i k a_0 b_2 x_1)$$

This is the class (9). If we go to the quotient space $\mathbb{T}^3$, then the action (10) on the fibres defines a gerbe of topological type $\alpha \smile \beta$ where β is a Cech cohomology class associated to the line bundle with Chern class equal to k and α is the Cech generator as in 2.1.

References

[1] Atiyah M. F.: *K*-Theory (W. A. Benjamin, Inc. 1967)

[2] Behrend K., Xu P.: J. Symplectic Geom. 9, 285-341 (2011)

[3] Bouwknegt P., Carey A., Mathai V., Murray M., Stevenson D.: Twisted K-theory and K-theory of bundle gerbes. Comm. Math. Phys. 228, 17 - 45 (2002)

[4] Berline N., Getzler E., Vergne M.: Heat Kernels and Dirac Operators (Springer-Verlag, Berlin 2004)

[5] Carey A., Mickelsson J., Murray M.: Index theory, gerbes, and Hamiltonian quantization. Comm. Math. Phys. 183, 707-722 (1997)

[6] Carey A., Mickelsson J., Murray M.: Bundle Gerbes Applied to Quantum Field Theory, Rev. Math. Phys. 12, 65-90 (2000)

[7] Douglas C. L.: On the twisted *K*-homology of simple Lie groups, Topology 45, 955-988 (2006)

[8] Freed D.S., Hopkins M.J., Teleman C.: (2011) Loop groups and twisted K-theory I, J. Topol. 4, 737-798 (2011)

[9] Freed D.S., Hopkins M.J., Teleman C.: Loop Groups and Twisted K-Theory II, arXiv:0511232v2 [math.AT] (2007)

[10] Guillemin V., Ginzburg V., Karshon Y.: Moment maps, cobordisms, and Hamiltonian group actions, Mathematical Surveys and Monographs, 98 (AMS, Providence, RI 2002)

[11] Harju A. J., Mickelsson J.: Twisted K-theory constructions in the case of a decomposable Dixmier-Douady class, arXiv:1208:4921 [math.KT] (2012)

[12] Nicolaescu L.: On the Space of Fredholm Operators, An. Sti. Univ. Iasi 53, 209-227 (2007)

[13] Lott J.: Higher-Degree Analogs of the Determinant Line Bundle, Commun. Math. Phys. 230, 41-69 (2002)

[14] Melrose M.P., Piazza P.: Families of Dirac Operators, Boundaries and the B-Calculus, J. Diff. Geom. 45, 99-180 (1997)

[15] Mickelsson J.: Gerbes, (twisted) K-theory, and the supersymmetric WZW model, in Wurzbacher T.: Infinite Dimensional Groups and Manifolds, IRMA Lectures in Mathematics and Theoretical Physics 5 (Walter de Gruyter 2004)

[16] Mickelsson J.: Gerbes and Quantum Field Theory, in: Encyclopedia of Mathematical Physics Vol 2, 539-546 (Elsevier 2006)

[17] Mickelsson J., Pellonpaa J-P.: Families Index Theorem in Supersymmetric WZW Model and Twisted K-Theory: The $SU(2)$ Case, Commun. Math. Phys. 271, 775-789 (2007)

[18] Ottesen J. T.: Infinite Dimensional Groups and Algebras in Quantum Physics (Springer-Verlag, Berlin 1995)

[19] Quillen D.: Superconnections and the Chern Character, Topology 24, 88-95 (1985)

[20] Tu J-L., Xu P., Laurent-Gengoux C.: Twisted K-Theory of Differentiable Stacks, Ann. Scient. Ec. Norm. Sup. 37, 841-910 (2004)